

New guidelines for testing “Deep eutectic solvents” toxicity and their effects on the environment and living beings

Javier Torregrosa-Crespo, Xavier Marset, Gabriela Guillena, Diego J. Ramón, Rosa María Martínez-Espinosa

PII: S0048-9697(19)35375-6  
DOI: <https://doi.org/10.1016/j.scitotenv.2019.135382>  
Reference: STOTEN 135382

To appear in: *Science of the Total Environment*

Received Date: 1 August 2019  
Revised Date: 8 October 2019  
Accepted Date: 3 November 2019



Please cite this article as: J. Torregrosa-Crespo, X. Marset, G. Guillena, D.J. Ramón, R. María Martínez-Espinosa, New guidelines for testing “Deep eutectic solvents” toxicity and their effects on the environment and living beings, *Science of the Total Environment* (2019), doi: <https://doi.org/10.1016/j.scitotenv.2019.135382>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**New guidelines for testing “Deep eutectic solvents” toxicity and their effects on the environment and living beings.**

Javier Torregrosa-Crespo<sup>1</sup>, Xavier Marset<sup>2</sup>, Gabriela Guillena<sup>2</sup>, Diego J. Ramón<sup>2</sup>, Rosa María Martínez-Espinosa\*<sup>1</sup>

<sup>1</sup>Departamento de Agroquímica y Bioquímica. División de Bioquímica y Biología Molecular. Facultad de Ciencias. Universidad de Alicante. Apdo. 99, E-03080-Alicante, Spain.

<sup>2</sup>Instituto de Síntesis Orgánica (ISO) and Departamento de Química Orgánica. Facultad de Ciencias. Universidad de Alicante. Apdo. 99, E-03080-Alicante, Spain.

\*Correspondence: rosa.martinez@ua.es; Tel.: +34965903400 ext. 1258; 8841

**New guidelines for testing “Deep eutectic solvents” toxicity and their effects on the environment and living beings.**

Javier Torregrosa-Crespo<sup>1</sup>, Xavier Marset<sup>2</sup>, Gabriela Guillena<sup>2</sup>, Diego J. Ramón<sup>2</sup>, Rosa María Martínez-Espinosa\*<sup>1</sup>

<sup>1</sup>Departamento de Agroquímica y Bioquímica. División de Bioquímica y Biología Molecular. Facultad de Ciencias. Universidad de Alicante. Apdo. 99, E-03080-Alicante, Spain.

<sup>2</sup>Instituto de Síntesis Orgánica (ISO) and Departamento de Química Orgánica. Facultad de Ciencias. Universidad de Alicante. Apdo. 99, E-03080-Alicante, Spain.

\*Correspondence: rosa.martinez@ua.es; Tel.: +34965903400 ext. 1258; 8841

**Highlights**

- DESs are promising compounds for organic chemistry and pharmacy.
- Despite their advantages as reaction media, there is controversy about their toxicity.
- Their toxicity was mainly assayed by antibiograms, which is not an accurate approach for DESs.
- New guidelines for DESs toxicity monitoring are presented.
- Protocols to test DES toxicity may lead to errors in assigning their eco-friendly nature.

**Abstract**

Deep eutectic solvents (DESs) were described at the beginning of this century as an alternative to ionic liquids (ILs) in green chemistry. Despite their obvious sustainable advantages as reaction media, there is still controversy about their potential toxicity. Most of the ecotoxicity assays done up to now involving DESs are based on antibiograms. This is not a good approach due to the high density and viscosity of most DESs already described. Additionally, antibiograms do not allow continuous monitoring of neither cellular growth nor changes on physicochemical parameters like culture acidification due to cellular growth or DESs metabolization. This work starts by displaying advantages and disadvantages of the DESs toxicity assays already reported. Then, using a new DES recently described and *Escherichia coli* as a model microorganism, liquid cultures with continuous monitoring of pH, temperature, shaking and optical density have been used, for the first time, to quantify potential toxicity of the DES as well as the degree of the cellular tolerance (in preadapted and non-preadapted cells). The results obtained show that this new DES is not toxic for *E. coli* at concentrations up to 300 mM and cellular preadaptation was crucial for the cells to grow. At concentrations between 300 mM and 450 mM, cells can tolerate this DES. Above 600 mM, the DES is toxic causing complete inhibition of growth. This toxicity is not only due to the chemical composition of the DES, but also due to the high acidification of the media caused by the DES hydrolysis during cellular growth. The consequences of sterilization procedures on the DES stability are also analysed into detail, finding that sterilization by autoclave promotes DES hydrolysis. From these results, new guidelines are proposed for further studies aiming to characterize and quantify DESs toxicity.

**Keywords:** Deep eutectic solvents, *Escherichia coli*, cellular toxicity, environmental toxicity.

## 1. Introduction

Deep eutectic solvents (DESs) are solvents generally formed by mixing a variety of quaternary ammonium salts with a Brønsted or Lewis acid compound in certain proportion. They were described for the first time at the beginning of this century as an alternative to ionic liquids (ILs) in green chemistry and catalysis (Abbott et al., 2004; Xu et al., 2017). Since that time around 800 papers have been published about DES (Date of access to PubMed and Scopus: 15 July 2019), involving subjects like: i) synthesis of new DES including their physicochemical characterisation (Marset et al., 2019; Zhang et al., 2012); ii) potential applications of DES as green solvent in biocatalysis (both, in presence or absence of enzymes and/or cells of biotechnological interest) (Gorke et al., 2008; Gutiérrez et al., 2010; Lindberg et al., 2010; Zhao et al., 2011; Lynam et al., 2017; Gotor-Fernández et al., 2019; Pätzold et al., 2019); iii) DES as ingredients for drugs synthesis, solubilization and delivery (Morrison et al., 2009; Pedro et al., 2019; Gutiérrez et al., 2019); iv) DES and/or NADES (Natural Deep Eutectic Solvents) as solvent for extraction of biomolecules like polysaccharides, proteins, alkaloids, phenolic compounds or other natural compounds like chitin, pigments, flavonoids,  $\kappa$ -carrageenan, dyes (Gu et al., 2014; Pena-Pereira et al., 2014; Zeng et al., 2014; García et al., 2016; Zhu et al., 2017; Zhuang et al., 2017; Saravana et al., 2018; Takla et al., 2018; Zdanowicz et al., 2018; Zhang et al., 2018; Huang et al., 2019) or v) DES as media for structural studies of nucleic acids (Mamajanov et al., 2010; Zhao et al., 2013a; Zhao et al., 2013b). More recently, DES-based molecularly imprinted polymers or DES-functionalised nanotubes have been developed and implemented at industrial scale for the removal of contaminants from wastewater or for the screening on antibiotics in biological fluids like milk (AlOmar et al., 2017; Tang et al., 2017; Chen et al., 2018).

Combinations of DES applications with other techniques or approaches such as nanoparticles/nanospheres and electrochemistry have also been reported (Nkuku and LeSuer, 2007; Abbott et al., 2009; Sheng et al., 2012). The main results described up to now reveal DESs as promising solvents not only for green chemical industries (Dai et al., 2013), but also for those related to cosmetics, food, pharmacy, biotechnology and biomedicine (Wagle et al., 2014; Cvjetko Bubalo et al., 2016; Li et al., 2016; Mbous et

al., 2017; Aydin et al., 2018; Jablonský et al., 2018; Procentese et al., 2018; Santos et al., 2018; Yoo et al., 2018).

In order to consider DESs as environmental friendly solvents, their real toxicity as well as their potential biodegradability must be established (Radošević et al., 2015). Studies regarding these facts have been performed to a lesser extent; mainly using prokaryotic microbes (Juneidi et al., 2016) and in some cases eukaryotic organisms (Radošević et al., 2016; Dai et al., 2014). Nevertheless, there is still controversy about their potential toxicity (Hayyan et al., 2013a). Whilst several reports support the theory that DESs are non-toxic, eco-friendly, biodegradable and benign solvents, others demonstrate the opposite (Hayyan et al., 2015; Hebert et al., 2016). In general terms, it has been assumed that they are not toxic and environmentally friendly because most of the DES components are naturally occurring, bio-renewable, biodegradable or can be bio-assimilated.

The standard protocols to study the cytotoxicity of new chemical components (such as the disk test) are not accurate in the case of DESs. The high viscosity, among other characteristics, makes the use of new methodologies to assess their biological toxicity essential. This microbiological study presents, for the first time, the continuous monitoring of various parameters (such as growth and pH) in liquid cultures with different concentrations of DES. This new protocol has revealed advantages in terms of biocompatibility, toxicity and tolerance.

## **2. Materials and Methods.**

### **2.1. Synthesis of DES.**

A recently described DES has been used in this study to optimize new protocols and guidelines for checking its toxicity. In brief, DES was synthesized using acetylcholine chloride (AcChCl):acetamide (1:2) as components. This was used as reaction medium to produce sulphonamides using a copper-catalysed process starting from triarylbiomethines,  $\text{Na}_2\text{S}_2\text{O}_5$  and nitro compounds (Marset et al., 2019).

### **2.2. Microbial strain, culture media and incubation conditions.**

The microorganism used in this study to test the toxicity, tolerance and the potential assimilation of the DES described here was *Escherichia coli* BL21 (DE3)

(Novagen). This microbe is a good example of a bacterial model organism extensively used in molecular biology, microbiology and genetic engineering due to its short generation time and large relevance in applied microbiology and biotechnology processes like protein overexpression (Kim et al., 2017).

The incubation experiments were conducted in 250 mL Erlenmeyers with 50 mL of Luria-Bertani medium (LB) following two strategies (Sezonov et al., 2007). On one hand, the initial pH of the culture media was adjusted to 7.0 using NaOH or HCl. Alternatively, the culture media were buffered with Tris-HCl 100 mM, adjusting the initial pH to 7.0. In all cases, pH was continuously monitored. The media was inoculated with 50  $\mu$ L of a pre-inoculum of cells previously grown in standard LB media until the stationary phase of growth or with 50  $\mu$ L of *E. coli* cells previously grown in LB media supplemented with DES at a lower concentration than the one tested in each assay. In all cases, the cells were grown at 37 °C with constant shaking (180 rpm). The growth of the bacterium was monitored by measuring the absorbance at 600 nm. All the physiological studies were done in triplicate.

In order to test the toxicity/tolerance of the DES, the media were supplemented with it, having final concentrations between 0 (control) and 750 mM. Likewise, the potential toxicity of the components that form this DES was also evaluated separately: acetylcholine chloride (AcChCl) and acetamide (MeCONH<sub>2</sub>). The concentrations used were 600 mM and 1.2 M, respectively, which correspond to those contained in the mixture of DES at 600 mM (maximum concentration at which cell growth was detected). DES, acetylcholine chloride and acetamide were sterilised by following two approaches: they were either added to the culture media before autoclaving them at 120 °C for 30 minutes; or the DES was sterilized by UV and acetylcholine chloride and acetamide were filtered through 0.22  $\mu$ m size filter and then added to the sterilised culture media.

The inhibition effect of the DES, AcChCl and MeCONH<sub>2</sub> was estimated by using the inhibition index (I) defined as:  $I = (A_0 - A) / A_0 * 100\%$  (Kommanee et al., 2012), where  $A_0$  and A are the absorbances at 600 nm for the cultures in the absence and presence of the DES, AcChCl and MeCONH<sub>2</sub>, respectively. Experiments were done by triplicate.



### 3. Results.

#### 3.1. Analysis of pH changes in the culture media during the cell growth in the presence of DES.

A few studies about the toxicity of DES using bacteria as model organisms suggest that pH decreases significantly in culture media supplemented with DES, however, pH has not been continuously monitored in these studies (Dai et al., 2014). For this reason, the first analysis introduced in this study was the continuous checking of pH in buffered and unbuffered culture media supplemented with DES, acetylcholine chloride (AcChCl) or acetamide (MeCONH<sub>2</sub>). The most dramatic pH depletion was detected in those unbuffered media in which DES or AcChCl were added to the culture media before autoclaving. Table 1 summarizes pH values at different time during the cell growth comparing buffered and unbuffered media. Even when the cultures were buffered, pH of the media containing DES or AcChCl decreased significantly (down to 4.4 in unbuffered DES-containing cultures and 4.5 in unbuffered and buffered AcChCl-containing cultures). The inoculation of the cultures and subsequent growth of the cells slightly increases the pH value as expected in the case of the controls and in the media containing MeCONH<sub>2</sub> (buffered and unbuffered).

[TABLE 1]

When the DES, AcChCl or MeCONH<sub>2</sub> were previously sterilised by UV or filtration respectively, and then added to the sterilised media, the changes of pH values were smaller (down to 5.3 as final pH in some of the cultures). In order to identify the magnitude of the contribution of autoclave and microbial metabolism to pH depletion, pH values were monitored in non-inoculated culture media containing two different concentrations of DES (150 mM and 600 mM). These cultures were incubated for the same period, under the same shaking and temperature conditions. After 24 hours of incubation, the final pH measured in sterile cultures with 600 mM of DES was 6.05 (Supplementary Table S1), which was much higher than the value for the same DES concentration inoculated with cells (Table 1).

From the results displayed in Table 1 and Supplementary Table S1, it is possible to conclude that the sterilization process based on autoclave may promote DES hydrolysis and probably reactions involving AcChCl, which contribute to pH acidification. DES hydrolysis and the consequent pH depletion also take place during the incubation, due to the culture media (Supplementary Table S1). *E. coli* growth and its metabolic activity is also involved in pH depletion.

### 3.2. Analysis of *E. coli* growth in the presence of different DES concentrations

To analyse *E. coli* tolerance and therefore the potential DES toxicity, cells were grown in buffered LB media supplemented with 600 mM DES and inoculated with *E. coli* cells grown in standard LB media (non-preadapted cells). This DES concentration was chosen based on the concentrations previously reported in the literature (Coleman et al., 2010). Cells were not able to grow under these conditions and the preliminary conclusion pointed out by the authors in this work was that 600 mM DES is toxic for this bacterium. It is important to highlight that microorganisms in general can induce specific reactions or pathways when they are exposed to compounds that are unusual in their natural environments or microcosmos in the laboratory. Considering this, several physiological studies were done to analyse the growth of pre-adapted *E. coli* cells. Thus, cells grown in the presence of 150 mM DES were used as inoculum for cultures containing 300 mM, etc. Figure 1 displays the growth curves of *E. coli* pre-adapted cells in LB containing different concentrations of DES.

#### [FIGURE 1]

The curves summarized in Figure 1 show two different patterns. The control curve shows the expected pattern in which a short Lag phase is followed by an exponential phase of growth (Lag phase) and then by a stationary phase of growth in which the maximum optical density (OD) observed was around 4. Shortly after 15 hours of incubation, the death phase starts. Curves from culture media containing 150 mM and 300 mM DES showed similar pattern and the final OD was close to 4. Regarding the cultures supplemented with 150 mM DES, two different curves can be

distinguished in Figure 1: one was obtained from culture media inoculated with cells from the standard culture (LB with no DES), and the other was measured from cultures in which cells used as inoculum were previously grown at 150 mM DES (preadapted cells). The differences found were clear; the culture inoculated with preadapted cells reduced its adaptation time (Lag phase) by two hours and cell duplication rate was increased (the slope of exponential phase of growth goes from 0.464 to 0.504). Therefore, better results in terms of cellular growth were obtained with preadapted cells thus suggesting that *E. coli* can deal with DES at least at concentrations between 0 and 300 mM. The cell inhibition rate remains below 6% in all these cases (Table 2).

In order to identify the maximum concentration of DES tolerated/assimilated by *E. coli*, the rest of the cultures with higher DES concentrations were always inoculated with cells pre-adapted to the concentration of DES immediately prior to the exposure. When the limit of 300 mM was exceeded, cellular adaptation started. At concentrations of 450 and 600 mM, the growth curves showed diauxic form and higher Lag phase. Diauxic curves have been previously reported from *E. coli* and many other bacteria (diphasic growth represented by two growth curves intervened by a short lag phase). These could be the result of: i) cellular growth usually using two different substrates at a different time (one of the substrates could be used preferentially and when it is exhausted, the second substrate is metabolised), or, ii) cellular growth affected by physicochemical parameters that can be changing during the incubation period. The maximum OD shown by the cultures with 450 mM DES was slightly higher (up to 4.5) than those reported from cultures with lower DES concentration. This profile could suggest that pre-adapted *E. coli* cells were able to grow nicely in the presence of 450 mM DES and may metabolise the DES or its derivatives when the DES is hydrolysed.

In the case of 600 mM, the percentage of cellular inhibition exceeded 30%, the Lag phase of growth increased significantly, and the final OD is half of the OD density reached by the other cultures. These results suggested that concentrations around 600 mM were toxic for the cells thus negatively affecting cellular growth and metabolic activities.

[TABLE 2]

The highest concentration tested in our experiments was 750 mM, at this concentration cell growth was not detected. Consequently, the DES concentrations above this value are toxic for *E. coli*.

### 3.3. Analysis of pH changes in culture media in the presence of the individual components of the DES: AcChCl and MeCONH<sub>2</sub> during *E. coli* growth

Regarding to the potential toxicity of the individual components of the DES here described (acetylcholine chloride – AcChCl – and acetamide – MeCONH<sub>2</sub> -), *E. coli* cells were grown in the presence of 600 mM DES (higher DES concentration allowing cellular growth) as well as in the presence of the two individual components at their final concentrations (600 mM AcChCl and 1.2 M MeCONH<sub>2</sub>).

#### [FIGURE 2]

The value of the Lag phase is closed in the three cultures (DES, AcChCl and MeCONH<sub>2</sub>; between 25 and 28 hours) but is higher than the Lag phase observed in control cultures. Final maximum OD in MeCONH<sub>2</sub>-culture is higher than in control cultures. However, final OD in DES and AcChCl cultures (around 2.5) is lower than in control cultures. DES is more toxic than its constituent elements separately, based on the higher inhibition percentage and the lower duplication tax (Supplementary Table S2). pH decrease was observed in both AcChCl-cultures and MeCONH<sub>2</sub>-cultures with 6.5 vs 7.4 at the end of the stationary phase of growth respectively. This depletion was higher in the presence of AcChCl, which correlates to the lower cellular growth detected in these cultures compared to MeCONH<sub>2</sub>-cultures. Comparing the cellular rates from the DES, AcChCl and MeCONH<sub>2</sub> cultures, it can be concluded that the individual components are less toxic than the DES (mixture of two components in 1:2 ratio) as has been previously reported for most of DESs described up to now.

### 4. Discussion.

New advances on the knowledge of applied organic chemistry, biomedicine, pharmacy, nanotechnologies and materials sciences are contributing to the improvement of several industrial/technical processes at global scale. However, there is a concern about the importance of assessing risks of environmental pollutants obtained in those processes (Nilsen et al., 2019; Vita et al., 2018). Thus, several recent works address new guidelines and approaches to optimize the assessment of the quite recently environmental risks in both, terrestrial and aquatic environments (Song et al., 2018; Zhang et al., 2019; Shiyuan et al. 2019). This work contributes to the knowledge of assessing risks of environmental pollutants, as it is the case of DESs as potential environmental contaminants. The search of literature focused on DESs using PubMed or Scopus as search engines revealed that around 800 papers have been published during the last 15 years (date of access 18 July 2019). The first studies were reported at the beginning of this century, but 90% of them have been mainly published during the last 5 years. Considering the recent high impact of these kind of solvents, it is worthy to highlight that only around 50 of these studies, analyse or discuss about potential toxicity on DESs (PubMed; “Deep eutectic solvents” AND “toxicity” as keywords). The study of DES toxicity is a quite new topic for research. Thus, the first approach for testing toxicity of choline chloride (ChCl) based DESs used two Gram positive bacteria *Bacillus subtilis* and *Staphylococcus aureus*, and two Gram negative bacteria *Escherichia coli* and *Pseudomonas aeruginosa*. The cytotoxic effect of choline chloride (ChCl) based DESs was tested using the *Artemia salina* leach at that time (Hayyan et al., 2013a). Shortly after this study, some others stated that DESs toxicological effects are different based on their components, and usually, the cytotoxicity of tested DESs was much higher than that of their individual components (Hayyan et al., 2013b).

More recent studies have tested the toxicity of choline chloride-based DESs, with glucose, glycerol and oxalic acid as hydrogen bond donors, using fish and human cell lines, phytotoxicity using wheat and biodegradability using wastewater microorganisms through closed bottle test (Radošević et al., 2015). The cytotoxic potential of ammonium-based DESs with four hydrogen bond donors has also been studied using a variety of cell lines *in vitro* and animal model *in vivo* (Hayyan et al., 2015; Chen et al., 2017; Macário et al., 2019). Other organisms chosen as model organisms for testing DESs toxicity have been the following: *Allium sativum* (plant) and

hydra (invertebrate) (Wen et al., 2015), several eukaryotic microorganisms like *Phanerochaete chrysosporium*, *Aspergillus niger*, *Lentinus tigrinus* and *Candida cylindracea* (Juneidi et al., 2016) and fish cell lines (Radošević et al., 2016)

Although the examined DESs showed in general less cytotoxicity compared with ionic liquids (Kudlak et al., 2015), not all DESs can be considered readily biodegradable or environmentally friendly (Wen et al., 2015). Besides, most of the test used up to now to monitor DESs toxicity, are based on antibiograms and cells are not usually preadapted. Considering that DESs are normally highly dense and viscous solvents, the approaches already used in most of the cases do not guarantee accurate assays in which DESs could diffuse and interact efficiently with the cells or microbes. Consequently, more investigation on this matter are required in order to provide specific and optimised protocols for testing DESs toxicity thus avoiding misunderstanding or wrong conclusions about potential negative effects of DESs.

A deep analysis of all studies reported up to now about the toxicity and biodegradability of DESs reveals that:

- A large proportion of DESs are considered as “biodegradable” because most the components forming DESs are natural products. However, in most of the reports, accurate experiments to test biodegradability have not been conducted.

- The terms biodegradation, assimilation, toxicity, tolerance and recycling are usually used in a confusing way in several studies based on DESs. Biodegradation involves degradation, transformation or accumulation of DESs or its derivatives within the cells. Assimilation involves the use of DESs as a source of nutrients for the growth of cells. Tolerance implies an increase of the biomass in a specific range of concentrations, but above the maximum concentration, cell growth and its metabolism become inhibited.

- In many of the experiments done with microbial cells to test the toxicity of DESs, parameters like rate of growth, time of incubation, OD at stationary phase of growth, pH of the media in unbuffered and buffered media are not continuously monitored. This is also true of some studies previously done using bacteria as model organism (Wen et al., 2015). As reported here, the addition of DES to the culture media promotes pH decrease. In addition to this, the process to sterilise the culture media or DES stock is also directly connected to pH decrease. This phenomenon was

also previously described in the literature (Dai et al., 2014). In our study, pH depletion due to DES hydrolysis or spontaneous reactions involving  $\text{MeCONH}_2$ , more significantly affects cellular proliferation and metabolic rates, than the DES or its singular components themselves.

- Toxicity can be measured by different approaches. In most of the reported works, toxicity is checked by using solid culture media in plates with disks in which DES is embedded (antibiogram-like test) (Zhao et al., 2015). This approach has an important limitation due to the high density and viscosity characterising most of the DESs described so far (Xu et al., 2017). These two properties restrict DESs diffusion from the disk to its surroundings, thus giving results that may not reflect real interaction between DESs and cells. Analysis of toxicity in liquid media minimizes this negative impact, especially in those cases in which DESs are particularly dense or show high viscosity.

- Biodegradability of DES (aerobic biodegradability) is usually evaluated using a closed bottle test, in which DESs are added to aerobic aqueous media inoculated with wastewater microorganisms, and the depletion of dissolved  $\text{O}_2$  is periodically determined (Coleman et al., 2010). The biodegradation test is normally considered as valid if a 60% ThOD is achieved within 14 d for reference compound, and chemicals are deemed to be readily biodegradable if they can reach a biodegradation level higher than 60% within a 28-d period (OECD, 1992). However, most of the microbial strains used in previous studies are microaerobic, anaerobic or can switch their metabolism from highly oxidic conditions to highly anoxic conditions. These aspects usually are not considered in the test of biodegradability done with DESs reported so far.

- Crossed reactions between DESs and the salts/nutrients of the culture media have not been considered in the toxicity studies. Hydrolysis of DESs could occur in liquid culture media thus promoting crossing reactions between the products of the DES hydrolysis and the salts, amino acids, carbohydrates, etc. within the culture media. Besides, recent studies demonstrate that the DES water content, as well as the formation of water shell layers are important in order to keep stable their nanostructure (Hammond et al., 2017; Ma et al., 2018; Kumari et al., 2018).

- None of the studies reported up to now have considered testing the preadaptation of cells to the DESs. Because of this, there is a loss of information in



terms of real capability of the cells to tolerate or even assimilate DESs. This study has demonstrated that under single assays in the presence of 600 mM DES the DES described here is toxic for the cells. However, when they were pre-adapted to this concentration, growth could be monitored. Consequently, we suggest including this parameter (cellular pre-adaptation) in further studies, in order to described more accurately the extent to which DESs are toxic to microbes.

## 5. Conclusions.

The new DES here tested is not toxic for *E. coli*, and maybe for other similar species, at concentrations between 0 and 300 mM. The DES used in this range of concentrations could be labelled as “environmentally friendly”. At concentrations between 300 mM and 450 mM, cells can tolerate the DES, even though cellular growth and metabolic activities are slightly affected by it (typical microbial growth curves change to diauxic –diphasic- or triauxic –triphasic- curves and Lag times are higher than those observe at lower DES concentrations). Above 600 mM, the DES is toxic causing complete inhibition of growth. This toxicity is not only due to the chemical composition of the DES, but also due to the high acidification of the media caused by DES hydrolysis. Thus, concentrations above 600 mM could allow its use as bactericide or even to explore potential inhibitory effect on tumoral cells. The assays described here have demonstrated that the type of culture used to test toxicity, preadaptation of the cells used as a model to check cytotoxicity or the methods used to sterilise DESs can significantly affect the results when evaluating the degree of toxicity of DESs. Therefore, new guidelines and recommendations must be implemented in order to get accurate results when describing DES toxicity: i) liquid culture media vs solid media or antibiograms; ii) preadapted cells/tissues/seeds vs. non preadapted cells/tissues/seeds; iii) continuous monitoring of OD, pH, temperature, radiation, etc. during the cellular incubation in the presence of DES vs. the absence of continuous monitoring of these parameters, iv) buffered culture media vs. unbuffered cultures and v) sterilization of DES by UV or filtration (0.22  $\mu$ m filters) vs. sterilization by autoclave.



More investigation is still required, and some interesting questions arise for the next future: Which are the molecular bases beyond DESs toxicity? Are there extremophilic organisms able to deal with high concentrations of DESs?

## References

Abbott, A.P., Boothby, D., Capper, G., Davies, D.L., Rasheed, R.K. 2004. Deep eutectic solvents formed between choline chloride and carboxylic acids: versatile alternatives to ionic liquids. *J. Am. Chem. Soc.* 126(29), 9142-9147. doi:10.1021/ja048266j

Abbott, A.P., El Ttaib, K., Frisch, G., McKenzie, K.J., Ryder, K.S. 2009. Electrodeposition of copper composites from deep eutectic solvents based on choline chloride. *Phys. Chem. Chem. Phys.* 11(21), 4269-4277. doi: 0.1039/b817881j

Alomar, M.K., Alsaadi, M.A., Jassam, T.M., Akib, S., Ali Hashim, M. 2017. Novel deep eutectic solvent-functionalized carbon nanotubes adsorbent for mercury removal from water. *J. Colloid. Interface. Sci.* 497, 413-421. doi:10.1016/j.jcis.2017.03.014

Aydin, F., Yilmaz, E., Soylak, M. 2018. Vortex assisted deep eutectic solvent (DES)-emulsification liquid-liquid microextraction of trace curcumin in food and herbal tea samples. *Food. Chem.* 243, 442-447. doi: 10.1016/j.foodchem.2017.09.154

Chen, J., Wang, Q., Liu, M., Zhang, L. 2017. The effect of deep eutectic solvent on the pharmacokinetics of salvianolic acid B in rats and its acute toxicity test. *J. Chromatogr. B. Analyt. Technol. Biomed. Life. Sci.* 1063, 60-66. doi: 10.1016/j.jchromb.2017.08.016

- Chen, J., Wang, Y., Wei, X., Xu, P., Xu, W., Ni, R., Meng, J. 2018. Magnetic solid-phase extraction for the removal of mercury from water with ternary hydrosulphonyl-based deep eutectic solvent modified magnetic graphene oxide. *Talanta*. 188, 454-462. doi: 10.1016/j.talanta.2018.06.016
- Coleman, D., Gathergood, N. 2010. Biodegradation studies of ionic liquids. *Chem. Soc. Rev.* 39(2), 600-637. doi:10.1039/b817717c
- Cvjetko Bubalo, M., Ćurko, N., Tomašević, M., Kovačević Ganić, K., Radojčić Redovniković, I. 2016. Green extraction of grape skin phenolics by using deep eutectic solvents. *Food. Chem.* 200, 159-166. doi: 10.1016/j.foodchem.2016.01.040
- Dai, Y., van Spronsen, J., Witkamp, G.J., Verpoorte, R., Choi, Y.H. 2013. Natural deep eutectic solvents as new potential media for green technology. *Anal. Chim. Acta.* 766, 61-68. doi:10.1016/j.aca.2012.12.013
- Dai, Y., Verpoorte, R., Choi, Y.H. 2014. Natural deep eutectic solvents providing enhanced stability of natural colorants from safflower (*Carthamus tinctorius*). *Food. Chem.* 159, 116-121. doi:10.1016/j.foodchem.2014.02.155
- García, A., Rodríguez-Juan, E., Rodríguez-Gutiérrez, G., Ríos, J.J., Fernández-Bolaños, J. 2016. Extraction of phenolic compounds from virgin olive oil by deep eutectic solvents (DESs). *Food. Chem.* 197(Pt A), 554-561. doi:10.1016/j.foodchem.2015.10.131
- Gorke, J.T., Sreenc, F., Kazlauskas, R.J. 2008. Hydrolase-catalyzed biotransformations in deep eutectic solvents. *Chem. Commun (Camb)*. 10, 1235-1237. doi: 10.1039/b716317g
- Gotor-Fernández, V., Paul, C.E. 2019. Deep eutectic solvents for redox biocatalysis. *J. Biotechnol.* 293, 24-35. doi:10.1016/j.jbiotec.2018.12.018

- 535 Gu, T., Zhang, M., Tan, T., Chen, J., Li, Z., Zhang, Q., Qiu, H. 2014. Deep eutectic  
536 solvents as novel extraction media for phenolic compounds from model oil. Chem.  
537 Commun (Camb). 50(79), 11746-11752. doi:10.1039/c4cc04661g  
538
- 539 Gutiérrez, A., Aparicio, S., Atilhan, M. 2019. Design of arginine-based therapeutic deep  
540 eutectic solvents as drug solubilization vehicles for active pharmaceutical ingredients.  
541 Phys. Chem. Chem. Phys. 21(20), 10621-10634. doi:10.1039/c9cp01408j  
542
- 543 Gutiérrez, M.C., Ferrer, M.L., Yuste, L., Rojo, F., del Monte, F. 2010. Bacteria  
544 incorporation in deep-eutectic solvents through freeze-drying. Angew. Chem. Int. Ed.  
545 Engl. 49(12), 2158-2162. doi:10.1002/anie.200905212  
546
- 547 Hammond, O.S., Bowron, D.T., Edler, K.J. 2017. The Effect of Water upon Deep Eutectic  
548 Solvent Nanostructure: An Unusual Transition from Ionic Mixture to Aqueous Solution.  
549 Angew Chem Int. Ed. Engl. 56(33), 9782-9785. doi: 10.1002/anie.201702486.  
550
- 551 Hayyan, M., Hashim, M.A., Hayyan, A., Al-Saadi, M.A., AlNashef, I.M., Mirghani, M.E.,  
552 Saheed, O.K. 2013a. Are deep eutectic solvents benign or toxic? Chemosphere. 90(7),  
553 2193-2195. doi:10.1016/j.chemosphere.2012.11.004  
554
- 555 Hayyan, M., Hashim, M.A., Al-Saadi, M.A., Hayyan, A., AlNashef, I.M., Mirghani, M.E.  
556 2013b. Assessment of cytotoxicity and toxicity for phosphonium-based deep eutectic  
557 solvents. Chemosphere. 93(2), 455-459. doi: 10.1016/j.chemosphere.2013.05.013  
558
- 559 Hayyan, M., Looi, C. Y., Hayyan, A., Wong, W. F. & Hashim, M. A. 2015. *In Vitro* and *In*  
560 *Vivo* toxicity profiling of ammonium-based deep eutectic solvents. PLoS One 10(2),  
561 e0117934/0117931-e0117934/0117918, doi:10.1371/journal.pone.0117934  
562
- 563 Hébert, M., Petiot, P., Benoit, E., Dansereau, J., Ahmad, T., Le Roch, A., Ottenwaelder,  
564 X., Gagnon, A. 2016. Synthesis of highly functionalized triarylbi-muthines by functional  
565 group manipulation and use in palladium- and copper-catalyzed arylation reactions. J.  
566 Org. Chem. 81(13), 5401-5416. doi:10.1021/acs.joc.6b00767

567

568 Huang, J., Guo, X., Xu, T., Fan, L., Zhou, X., Wu, S. 2019. Ionic deep eutectic solvents for  
 569 the extraction and separation of natural products. J. Chromatogr. A. 1598, 1-19.  
 570 doi:10.1016/j.chroma.2019.03.046

571

572 Jablonský, M., Škulcová, A., Malvis, A., Šima, J. 2018. Extraction of value-added  
 573 components from food industry based and agro-forest biowastes by deep eutectic  
 574 solvents. J. Biotechnol. 282, 46-66. doi: 10.1016/j.jbiotec.2018.06.349

575

576 Juneidi, I., Hayyan, M., Mohd, Ali O. 2016. Toxicity profile of choline chloride-based  
 577 deep eutectic solvents for fungi and *Cyprinus carpio* fish. Environ. Sci. Pollut. Res. Int.  
 578 23(8), 7648-7659. doi: 10.1007/s11356-015-6003-4

579

580 Kim, S., Jeong, H., Kim, E.Y., Kim, J.F., Lee, S.Y., Yoon, S.H. 2017. Genomic and  
 581 transcriptomic landscape of *Escherichia coli* BL21(DE3). Nucleic Acids. Res. 45(9), 5285-  
 582 5293. doi:10.1093/nar/gkx228

583

584 Kommanee, J., Preecharram, S., Daduang, S., Temsiripong, Y., Dhiravisit, A., Yamada, Y.,  
 585 Thammasirirak, S. 2012. Antibacterial activity of plasma from crocodile (*Crocodylus*  
 586 *siamensis*) against pathogenic bacteria. Ann. Clin. Microbiol. Antimicrob. 11:22.  
 587 doi:10.1186/1476-0711-11-22

588

589 Kudłak, B., Owczarek, K., Namieśnik, J. 2015. Selected issues related to the toxicity of  
 590 ionic liquids and deep eutectic solvents-a review. Environ. Sci. Pollut. Res. Int. 22(16),  
 591 11975-11992. doi: 10.1007/s11356-015-4794-y

592

593 Kumari, P., Shobhna, Kaur, S, Kashyap, H.K. 2018. Influence of Hydration on the  
 594 Structure of Reline Deep Eutectic Solvent: A Molecular Dynamics Study. ACS Omega.  
 595 3(11), 15246-15255. doi: 10.1021/acsomega.8b02447.

596

597 Li, X., Row, K.H. 2016. Development of deep eutectic solvents applied in extraction and  
 598 separation. J. Sep. Sci. 39(18), 3505-3520. doi: 10.1002/jssc.201600633

599

600 Lindberg, D., de la Fuente Revenga, M., Widersten, M. 2010. Deep eutectic solvents  
 601 (DESs) are viable cosolvents for enzyme-catalyzed epoxide hydrolysis. *J. Biotechnol.*  
 602 147(3-4), 169-171. doi:10.1016/j.jbiotec.2010.04.011

603

604 Lynam, J.G., Kumar, N., Wong, M.J. 2017. Deep eutectic solvents' ability to solubilize  
 605 lignin, cellulose, and hemicellulose; thermal stability; and density. *Bioresour. Technol.*  
 606 238, 684-689. doi:10.1016/j.biortech.2017.04.079

607

608 Ma, C., Laaksonen, A. , Liu, C., Lu, X. , Ji, X. 2018. The peculiar effect of water on ionic  
 609 liquids and deep eutectic solvents. *Chem. Soc. Rev.* 47(23), 8685-8720. doi:  
 610 10.1039/c8cs00325d.

611

612 Macário, I.P.E., Oliveira, H., Menezes, A.C., Ventura, S.P.M., Pereira, J.L., Gonçalves,  
 613 A.M.M., Coutinho, J.A.P., Gonçalves, F.J.M. 2019. Cytotoxicity profiling of deep eutectic  
 614 solvents to human skin cells. *Sci Rep.* 9(1):3932. doi: 10.1038/s41598-019-39910-y

615

616 Mamajanov, I., Engelhart, A.E., Bean, H.D., Hud, N.V. 2010. DNA and RNA in anhydrous  
 617 media: duplex, triplex, and G-quadruplex secondary structures in a deep eutectic  
 618 solvent. *Angew. Chem. Int. Ed. Engl.* 49(36), 6310-6314. doi:10.1002/anie.201001561

619

620 Marset, X., Torregrosa-Crespo, J., Martínez-Espinosa, R.M., Guillena, G., Ramón, D.J.  
 621 2019. Multicomponent synthesis of sulfonamides from triarylbiurathines, nitro  
 622 compounds and sodium metabisulfite in deep eutectic solvents. *Green Chem.* 21,  
 623 4127-4132. doi:10.1039/c9gc01541h

624

625 Mbous, Y.P., Hayyan, M., Hayyan, A., Wong, W.F., Hashim, M.A., Looi, C.Y. 2017.  
 626 Applications of deep eutectic solvents in biotechnology and bioengineering-Promises  
 627 and challenges. *Biotechnol. Adv.* 35(2), 105-134. doi:  
 628 10.1016/j.biotechadv.2016.11.006

629

- Morrison, H.G., Sun, C.C., Neervannan, S. 2009. Characterization of thermal behavior of deep eutectic solvents and their potential as drug solubilization vehicles. *Int. J. Pharm.* 378(1-2), 136-139. doi:10.1016/j.ijpharm.2009.05.039
- Nilsen, E., Smalling, K.L., Ahrens, L., Gros, M., Miglioranza, K.S.B., Picó, Y., Schoenfuss, H.L. 2019. Critical review: Grand challenges in assessing the adverse effects of contaminants of emerging concern on aquatic food webs. *Environ. Toxicol. Chem.* 38(1), 46-60. doi: 10.1002/etc.4290.
- Nkuku, C.A., LeSuer, R.J. 2007. Electrochemistry in deep eutectic solvents. *J. Phys. Chem. B.* 111(46), 13271-13277. doi:10.1021/jp075794j
- OECD. 1992. Guideline for testing of chemicals, adopted by the Council on 17th July 1992 Ready Biodegradability (<https://www.oecd.org/chemicalsafety/risk-assessment/1948209.pdf>)
- Pätzold, M., Siebenhaller, S., Kara, S., Liese, A., Syldatck, C., Holtmann, D. 2019. Deep eutectic solvents as efficient solvents in biocatalysis. *Trends. Biotechnol.* pii:S0167-7799(19)30052-6. doi:10.1016./j.tibtech.2019.03.007
- Pedro, S.N., Freire, M.G., Freire, C.S.R., Silvestre, A.J.D. 2019. Deep eutectic solvents comprising active pharmaceutical ingredients in the development of drug delivery systems. *Expert. Opin. Drug. Deliv.* 16(5), 497-506. doi:10.1080/17425247.2019.1604680.
- Pena-Pereira, F., Namiesnik, J. 2014. Ionic liquids and deep eutectic mixtures: sustainable solvents for extraction processes. *Chem. Sus. Chem.* 7(7), 1784-1800. doi: 10.1002/cssc.201301192
- Procentese, A., Raganati, F., Olivieri, G., Russo, M.E., Rehmann, L., Marzocchella, A. 2018. Deep eutectic solvents pretreatment of agro-industrial food waste. *Biotechnol. Biofuels.* 11:37. doi: 10.1186/s13068-018-1034-y

662

663 Radošević, K., Bubalo, M.C., Srček, V.G., Grgas, D., Dragičević, T.L., Redovniković, I.R.  
 664 2015. Evaluation of toxicity and biodegradability of choline chloride based deep  
 665 eutectic solvents. *Ecotoxicol. Environ. Saf.* 112, 46-53.  
 666 doi:10.1016/j.ecoenv.2014.09.034

667

668 Radošević, K., Železnjak, J., Cvjetko Bubalo, M., Radojčić Redovniković, I., Slivac, I.,  
 669 Gaurina Srček V. 2016. Comparative *in vitro* study of cholinium-based ionic liquids and  
 670 deep eutectic solvents toward fish cell line. *Ecotoxicol. Environ. Saf.* 131, 30-36.  
 671 doi:10.1016/j.ecoenv.2016.05.005

672

673 Santos, F., P. S. Leitão, M.I, C. Duarte, A.R. 2018. Properties of therapeutic deep  
 674 eutectic solvents of L-arginine and ethambutol for tuberculosis treatment. *Molecules*.  
 675 24(1), pii: E55. doi: 10.3390/molecules24010055

676

677 Saravana, P.S., Ho, T.C., Chae, S.J., Cho, Y.J., Park, J.S., Lee, H.J., Chun, B.S. 2018. Deep  
 678 eutectic solvent-based extraction and fabrication of chitin films from crustacean waste.  
 679 *Carbohydr. Polym.* 195, 622-630. doi:10.1016/j.carbpol.2018.05.018

680

681 Sezonov, G., Joseleau-Petit, D., D'Ari, R. 2007. *Escherichia coli* physiology in Luria-  
 682 Bertani broth. *J. Bacteriol.* 189(23), 8746-8749. doi:10.1128/JB.01368-07

683

684 Sheng, Q., Liu, R., Zheng, J. 2012. Prussian blue nanospheres synthesized in deep  
 685 eutectic solvents. *Nanoscale*. 4(21), 6880-6886. doi:10.1039/c2nr31830j.

686

687 Song, B., Xu, P., Zeng, G., Gong, J., Wang, X., Yan, J., Wang, S., Zhang, P., Cao, W., Ye, S.  
 688 2018. Modeling the transport of sodium dodecyl benzene sulfonate in riverine  
 689 sediment in the presence of multi-walled carbon nanotubes. *Water Res.* 129, 20-28.  
 690 doi: 10.1016/j.watres.2017.11.003.

691

692 Takla, S.S., Shawky, E., Hammada, H.M., Darwish, F.A. 2018. Green techniques in  
 693 comparison to conventional ones in the extraction of Amaryllidaceae alkaloids: best

solvents selection and parameters optimization. *J. Chromatogr. A.* 1567, 99-110.  
doi:10.1016/j.chroma.2018.07.009

Tang, W., Gao, F., Duan, Y., Zhu, T., Ho Row, K. 2017. Exploration of deep eutectic solvents-based molecularly imprinted polymers as solid-phase extraction sorbents for screening chloramphenicol in milk. *J. Chromatogr. Sci.* 55(6), 654-661.  
doi:10.1093/chromsci/bmx011

Vita, N.A., Brohem, C.A., Canavez, A.D.P.M., Oliveira, C.F.S., Kruger, O., Lorencini, M., Carvalho, C.M. 2018. Parameters for assessing the aquatic environmental impact of cosmetic products. *Toxicol Lett.* 287, 70-82. doi: 10.1016/j.toxlet.2018.01.015.

Wagle, D.V., Zhao, H., Baker, G.A. 2014. Deep eutectic solvents: sustainable media for nanoscale and functional materials. *Acc. Chem. Res.* 47(8), 2299-2308.  
doi:10.1021/ar5000488

Wen, Q., Chen, J.X., Tang, Y.L., Wang, J., Yang, Z. 2015. Assessing the toxicity and biodegradability of deep eutectic solvents. *Chemosphere.* 132, 63-69.  
doi:10.1016/j.chemosphere.2015.02.061

Xu, P., Zheng, G.W., Zong, M.H., Li, N., Lou, W.Y. 2017. Recent progress on deep eutectic solvents in biocatalysis. *Bioresources and bioprocessing.* 4:34.  
doi:10.1186/s40643-017-0165-5

Yoo, D.E., Jeong, K.M., Han, S.Y., Kim, E.M., Jin, Y., Lee, J. 2018. Deep eutectic solvent-based valorization of spent coffee grounds. *Food. Chem.* 255, 357-364. doi: 10.1016/j.foodchem.2018.02.096

Zdanowicz, M., Wilpiszewska, K., Spychaj, T. 2018. Deep eutectic solvents for polysaccharides processing: a review. *Carbohydr. Polym.* 200, 361-380.  
doi:10.1016/j.carbpol.2018.07.078



- Zeng, Q., Wang, Y., Huang, Y., Ding, X., Chen, J., Xu, K. 2014. Deep eutectic solvents as novel extraction media for protein partitioning. *Analyst*. 139(10), 2565-2573. doi:10.1039/c3an02235h
- Zhang, H., Wang, Y., Zhou, Y., Chen, J., Wei, X., Xu, P. 2018. Aqueous biphasic systems formed by deep eutectic solvent and new-type salts for the high-performance extraction of pigments. *Talanta*. 181, 210-216. doi:10.1016/j.talanta.2018.01.014
- Zhang, M., Bradford, S.A., Šimůnek, J., Vereecken, H., Klumpp, E. 2019. Co-transport of multi-walled carbon nanotubes and sodium dodecylbenzenesulfonate in chemically heterogeneous porous media. *Environ. Pollut.* 247, 907-916. doi:10.1016/j.envpol.2019.01.106.
- Zhang, Q., De Oliveira Vigier, K., Royer, S., Jérôme, F. 2012. Deep eutectic solvents: syntheses, properties and applications. *Chem. Soc. Rev.* 41(21), 7108-7146. doi:10.1039/c2cs35178a
- Zhao, H., Baker, G.A., Holmes, S. 2011. Protease activation in glycerol-based deep eutectic solvents. *J. Mol. Catal. B. Enzym.* 72(3-4), 163-167. doi:10.1016/j.molcatb.2011.05.015
- Zhao, C., Qu, X. 2013a. Recent progress in G-quadruplex DNA in deep eutectic solvent. *Methods*. 64(1), 52-58. doi:10.1016/j.ymeth.2013.04.017
- Zhao, C., Ren, J., Qu, X. 2013b. G-quadruplexes form ultrastable parallel structures in deep eutectic solvent. *Langmuir*. 29(4), 1183-1191. doi:10.1021/la3043186
- Zhao, B.Y., Xu, P., Yang, F.X., Wu, H., Zong, M.H., Lou, W.Y. 2015. Biocompatible deep eutectic solvents based on choline chloride: characterization and application to the

extraction of rutin from *Sophora japonica*. ACS Sustainable Chem. Eng. 3(11), 2746-2755. doi:10.1021/acssuschemeng.5b00619

Zhou, S., Zheng, Z.; Mei, T.; Wang, X. 2019. Structural design and material preparation of carbon-based electrodes for high-performance lithium storage systems. Carbon, 144, 1-7.

Zhu, S., Liu, D., Zhu, X., Su, A., Zhang, H. 2017. Extraction of illegal dyes from red chili peppers with cholinium-based deep eutectic solvents. J. Anal. Methods. Chem. 2017, 2753752. doi: 10.1155/2017/2753752

Zhuang, B., Dou, L.L., Li, P., Liu, E.H. 2017. Deep eutectic solvents as green media for extraction of flavonoid glycosides and aglycones from *Platycladi Cacumen*. J. Pharm. Biomed. Anal. 134, 214-219. doi:10.1016/j.jpba.2016.11.049

**Acknowledgements**

This work was supported by the University of Alicante (VIGROB-173 and VIGROB-309), the Spanish Ministerio de Economía, Industria y Competitividad (CTQ2015-66624-P, PGC2018-096616-B-I00 and RTI2018-099860-B-I00). X.M. and J.T.C. thank Generalitat Valenciana (ACIF/2016/057 and ACIF/2016/077) for their fellowships. Thanks to T. Coyle for English checking of the manuscript.

**Author contributions**

R.M.M., D. J. R. and G. G. directed and supervised the project. X. M. designed and performed the synthetic experiments, characterized all products and performed the DES physicochemical characterization. J.T.C. and R.M.M. designed and performed the experiments related to the biological evaluation of the DES. All the authors prepared the manuscript and they have agreed to the content of it.

**Conflict of interest**

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.


**Table 1:** Average of pH values from three biological replicates of buffered and unbuffered culture media during the cell growth. (\*) means “buffered”. AcChCl corresponds to acetylcholine chloride and MeCONH<sub>2</sub> to acetamide. Final DES concentration was 600 mM. When AcChCl and MeCONH<sub>2</sub> were tested separately, their final concentrations were those contained in the mixture of DES at 600 mM (see materials and methods). C corresponds to control media in which LB is not supplemented neither DES nor AcChCl or MeCONH<sub>2</sub>. In all cases, DES, AcChCl and MeCONH<sub>2</sub> were added to the culture media before autoclaving.

Time (h of incubation)	C	C*	DES	DES*	AcChCl	AcChCl*	MeCONH <sub>2</sub>	MeCONH <sub>2</sub> *
0 before autoclave	7.0	7.0	6.9	6.9	6.9	6.9	6.9	7.0
0 after autoclave and before cells inoculation	7.2	7.0	4.5	4.7	4.5	4.7	6.6	7.0
5.0	7.5	6.8	4.4	4.6	4.5	4.5	6.4	6.9
10.3	8.0	7.1	4.5	4.6	4.4	4.6	6.9	6.8
23.0	8.5	8.1	4.5	4.7	4.4	4.6	8.4	7.7
48.0	8.4	8.1	4.5	4.8	4.5	4.7	8.5	7.9


**Table 2:** Summary of parameters regarding cellular growth and percentage of growth inhibition in culture media supplemented with different concentrations of the DES. The DES was sterilised by UV and then added to sterilised cultures. \* Indicates that cells used as inoculum were not pre-adapted.

[DES] (mM)	SLOPE ( $\Delta$ Abs/hour) (Rate of cellular duplication)	R <sup>2</sup>	I (%) (Percentage of growth inhibition)	Lag time (hours)	Average of pH values during incubation	Average of minimal pH value detected during incubation
0	0.67	0.99	0	3.75	6.87	6.77

150*	0.46	0.99	4.76	7.50	6.65	6.61
150	0.50	0.97	5.57	5.50	6.53	6.53
300	0.40	0.99	0.00	2.00	6.72	6.63
450	0.43	0.97	0.00	28.50	6.60	6.10
600	0.53	0.94	32.57	28.50	5.80	5.80
750	0	0	98.89	-	-	5.77



**Figure 1:** *E. coli* growth curves. Cells were grown in LB media supplemented with different concentrations of DES. Legend of colours: Orange = control (LB media without DES); Yellow = LB + 150 mM DES inoculated with pre-adapted cells; Grey = LB + 150 mM DES inoculated with non-pre-adapted cells; light blue = LB + 300 mM DES and preadapted cells; green = LB + 450 mM DES and preadapted cells; dark blue = LB + 600 mM DES and preadapted cells; dark red = LB + 750 mM DES and preadapted cells. For each curve, the dots correspond to the average of three values obtained from three biological replicates, standard deviation of these values was between  $\pm 0.01$  and  $\pm 0.16$ .



**Figure 2:** *E. coli* growth curves. Cells were grown in buffered LB media supplemented with 600 mM DES, 600 mM AcChCl or 1.2 M MeCONH<sub>2</sub>. DES was sterilised by UV and then added to sterilised media. AcChCl and MeCONH<sub>2</sub> were sterilised by filtration and then added to sterilised media. Legend of colours: Blue = control (LB media without DES); Purple = LB + 1.2 MeCONH<sub>2</sub>; Green = LB + 600 mM AcChCl; Red = LB + 600 mM DES. In all cases, cultures were inoculated with pre-adapted cells. For each curve, the dots correspond to the average of three values obtained from three biological replicates, standard deviations were between  $\pm 0.01$  and  $\pm 0.2$ .

#### SUPPLEMENTARY MATERIALS:

**Supplementary Table 1:** pH values in buffered media supplemented with 150 mM or 600 mM DES. DES was sterilised by UV and then added to sterilised cultures. These cultures were not inoculated with cells. SD values were  $\pm 0.1$  in all cases.

Time (hours of incubation)	150 mM DES	600 mM DES
0	7.00	7.00
2	6.80	6.90
6	6.65	6.75
10	6.52	6.35
15	6.41	6.25
24	6.27	6.05
48	5.65	5.30

**Supplementary Table 2:** summary of parameters related to cellular growth and percentage of growth inhibition in culture media supplemented with DES (600 Mm), acetylcholine chloride (AC) and acetamide (AT). DES was sterilized by UV while AC and AT were filtered using a 0,22-micron filter and then added to sterilized culture media.

Culture	SLOPE ( $\Delta$ Abs/hour)	R <sup>2</sup>	I (%)	Lag time	Average of	Average of
---------	----------------------------	----------------	-------	----------	------------	------------

	(Rate of cellular duplication)		(Percentage of growth inhibition)	(hours)	pH values during incubation	minimal pH value detected during incubation
Control	0.58	0.99	0.00	5.50	6.77	6.77
DES (600 mM)	0.36	0.93	21.44	5.50	6.34	5.87
AC	0.43	0.93	22.21	28.50	7.15	5.81
AT	0.34	0.98	0.00	28.50	6.10	6.10

### Highlights

- DESs are promising compounds for organic chemistry and pharmacy.
- Despite their advantages as reaction media, there is controversy about their toxicity.
- Their toxicity was mainly assayed by antibiograms, which is not an accurate approach for DESs.
- New guidelines for DESs toxicity monitoring are presented.
- Protocols to test DES toxicity may lead to errors in assigning their eco-friendly nature.

### Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.